ACTIFLORE® BO213

Selected Active Dry Yeasts (SADY), non-GMO, for oenological use.

Suitable for the preparation of products intended for direct human consumption, in the scope of regulated use in oenology.

Complies with Commission Regulation (EC) No. 2019/934.

SPECIFICATIONS AND OENOLOGICAL APPLICATIONS

Saccharomyces cerevisiae specially recommended for restarting fermentation. Fructose-fermenting strain with excellent resistance to alcohol and to low temperatures. ACTIFLORE® BO213 is also recommended for its aromatic neutrality (good respect for varietal character).

FERMENTATION CHARACTERISTICS

- Fructose-fermenting character.
- Alcohol tolerance: up to 18% vol.
- Fermentation temperature: 10 32°C (50 90°F).
- · Low nitrogen requirement.

AROMATIC CHARACTERISTICS

• Aromatically neutral (respects varietal character).

PHYSICAL CHARACTERISTICS

Vacuum-packed dehydrated yeasts

Appearance granules

CHEMICAL AND MICROBIOLOGICAL ANALYSIS

Humidity (%)
Activ Dry Yeast (ADY) (CFU/g) $\geq 1.2.10^{10}$
Lactic acid bacteria (CFU/g)< 10 st
Acetic acid bacteria (CFU/g) < 10 ⁴
Yeasts of a genus other than Saccharomyces (CFU/g) $< 10^{\circ}$
Yeasts of a different species or strain (%)< 5
Coliforms (CFU/g)< 10 ²
Moulds (CFU/g) < 10 ³

Staphylococcus (/g)none
Salmonella(/25 g) none
E. coli (/g)none
Lead (ppm)< 2
Arsenic (ppm) < 3
Mercury (ppm)<
Cadmium (ppm) < `



PROTOCOL FOR USE

OENOLOGICAL CONDITIONS

- Add the yeast as soon as possible after filling the tank.
- Respect the prescribed dose to ensure a good yeast implantation, even in case of abundance of indigenous yeasts.
- The temperature, choice of yeast strain, rehydration procedure and cellar hygiene are also essential for successful establishment of the population.

DOSAGE

- 20 30 g/hL (200 300 ppm) (depending on the must characteristics and vinification conditions).
- 30 50 g/hL (300 500 ppm) in case of stuck fermentation.

IMPLEMENTATION

- Carefully follow the yeast rehydration protocol.
- Avoid temperature differences greater than 10°C (50°F) between the must and the rehydrated yeast at the time of inoculation. The total yeast preparation time must not exceed 45 minutes.

In case of particularly difficult fermentation conditions (very low temperature, highly clarified must, very high potential alcohol) and/or to increase the aromatic potential of the wine, use SUPERSTART® or SUPERSTART®BLANC in the rehydration water.

STORAGE RECOMMENDATION

- Store above ground level in a dry area not liable to impart odours. Ensuring stock is kept at a moderate temperature, in its original, unopened packaging.
- Optimal date of use: 4 years.

PACKAGING

500 g vacuum bag. 10 kg box.10 kg bag.

